



Diffraction at the Tevatron



XXXIII International Conference on High Energy Physics
Moscow :: 28 July 2006



Andrew Hamilton
Université de Genève
for the CDF and DØ Collaborations

Introduction



introduction

outline

DØ plans

CDF inclusive

intro

results

CDF exclusive

intro



e^+e^-

$\gamma\gamma$

dijet

summary

Diffraction: An exchange with the quantum numbers of the vacuum.



Goals of Diffractive Program at the Tevatron:

- Understand the nature of diffractive exchange
- Test the feasibility of diffraction as a tool to search for new physics at the LHC

Outline



introduction

outline

DØ plans

CDF inclusive

intro

results

CDF exclusive

intro

e^+e^-

$\gamma\gamma$

dijet

summary

Diffraction at DØ:

- physics plans

Inclusive Diffraction at CDF:

- ratio of SD/ND dijets
- Q^2 dependence of t in SD dijets

Exclusive Production at CDF:

- two mechanisms (*QED and QCD*)
- searches for QED mediated e^+e^-
- searches for QCD mediated $\gamma\gamma$ and dijets

Summary

Diffractive Analyses in Progress at DØ:

introduction

outline

DØ plans

CDF inclusive

intro

results

CDF exclusive

intro

e^+e^-


$\gamma\gamma$

dijet


summary

- Double Pomeron Exchange (DPE) + Jets
- Diffractive Structure Function
- Diffractive Heavy Flavor
- Exclusive Production
- **Diffractive Z**
- Diffractive W
- Inclusive DPE

Di-Muon Invariant Mass - $0.02 < \xi_{pbar} < 0.1$



- introduction
- outline
- DØ plans
- CDF inclusive
 - intro
 - results
- CDF exclusive
 - intro
 - e^+e^-
 - $\gamma\gamma$
 - dijet
- summary



Use high p_T jets as a probe to measure **Diffraction Structure Function, F_{jj}^D**

Experimental Determination of F_{jj}^D

$$R(x_{Bj}) \text{ of } \frac{\sigma_{jj}(SD)}{\sigma_{jj}(ND)} = \frac{F_{jj}^D(x_{Bj}, Q^2)}{F_{jj}(x_{Bj}, Q^2)} \text{ (LO QCD)}$$


↑
Data

↑
Known Proton PDF

$$\beta = P_{parton} / P_{Pomeron}$$


$$\xi = P_{pomeron} / P_{proton}$$

$$t = (P_i - P_f)^2$$




- No Q^2 dependence
($100 < Q^2 < 10000 \text{ GeV}^2$)
- The Pomeron evolves like the proton


- Slope of $d\sigma/dt$ is independent of Q^2 in SD dijets



Exclusive Production: QED or QCD mediated



- measure exclusive QED e^+e^- and $\mu^+\mu^-$ production at Tevatron to test feasibility of Luminosity measurement at LHC



- measure exclusive QCD dijet and $\gamma\gamma$ production at Tevatron to constrain predictions on exclusive Higgs at LHC

introduction

outline

DØ plans

CDF inclusive

intro

results

CDF exclusive

intro

e^+e^-



$\gamma\gamma$

dijet

summary

Exclusive e^+e^- events are selected by:

- reconstructing the e^+e^-
- requiring that there is no other activity in $|\eta| < 7.4$
- *protons are not tagged*





16 events similar are found.



QED Mediated e^+e^- Production




introduction
outline
DØ plans
CDF inclusive
intro
results
CDF exclusive
intro
 e^+e^-
 $\gamma\gamma$
dijet
summary



Background estimate: $2.1^{+0.7}_{-0.3}$

$$\sigma_{\text{exp}} = 1.6^{+0.5}_{-0.3} (\text{stat}) \pm 0.3 (\text{sys}) \text{ pb}$$

$$\sigma_{\text{LPAIR}} = 1.711 \pm 0.008 \text{ pb}$$



introduction

outline

DØ plans

CDF inclusive

intro

results

CDF exclusive

intro

e^+e^-



$\gamma\gamma$

dijet

summary

Exclusive $\gamma\gamma$ events:

- selected in the same way as e^+e^- (except tracking)
- agreement of exclusive e^+e^- cross section gives confidence in analysis methodology



3 events are found.

1^{+3}_{-1} events are predicted from ExHuME MC


Monk & Pilkington. hep-ph/0502077

Background estimate is not yet complete



Strategy:

- select DPE dijets: $\bar{p} + p \rightarrow \bar{p} + X (\geq 2\text{jets} + \dots) + \text{gap}$
- examine the *dijet mass fraction* R_{jj}

$$R_{jj} = \frac{M_{jj}}{M_X}$$




introduction
outline
DØ plans
CDF inclusive
intro
results
CDF exclusive
intro
 e^+e^-
 $\gamma\gamma$
dijet
summary




Another Observable:

- exclusive gg → qq is suppressed by $J_Z=0$ rule
- exploit this by looking at ***fraction of heavy flavor (b/c) jets*** in dijet data as a function of R_{jj}

b/c jets suppressed at high R_{jj}



the two observables agree






QCD Mediated Dijet Production



introduction
outline
DØ plans
CDF inclusive
intro
results
CDF exclusive
intro
e⁺e⁻
γγ
dijet
summary

CDF Run II Preliminary




How does the cross section compare?

The data favors ExHuME MC

Ratio of exclusive to inclusive DPE cross sections is flat.

CDF Run II Preliminary



Summary



introduction

outline

DØ plans

CDF inclusive

intro

results

CDF exclusive

intro

e^+e^-

$\gamma\gamma$

dijet

summary

Diffraction at DØ:

- many analyses in progress

Inclusive Diffraction at CDF:

- Q^2 dependence of Diffractive Structure Function
- slope of SD dijet $d\sigma/dt$ is independent of Q^2

Exclusive Production at CDF:

- observed e^+e^- production via QED exchange
- possible evidence for $\gamma\gamma$ production via QCD exchange
- observed dijet production via QCD exchange
- KMR / ExHuME predictions are consistent with data
- see previous talk for impact KMR of predictions on the LHC



Backup Slides



- introduction
- outline
- DØ plans
- CDF inclusive
 - intro
 - results
- CDF exclusive
 - intro
 - e^+e^-
 - $\gamma\gamma$
 - dijet
- summary

Backup Slides



Tevatron Luminosity



introduction

outline

DØ plans

CDF inclusive

intro

results

CDF exclusive


intro

e^+e^-



$\gamma\gamma$

dijet

summary



- introduction
- outline
- DØ plans
- CDF inclusive
 - intro
 - results
- CDF exclusive
 - intro
 - e^+e^-
 - $\gamma\gamma$
 - dijet
- summary





ee Numbers



introduction

outline

DØ plans

CDF inclusive

intro

results

CDF exclusive

intro

e^+e^-

$\gamma\gamma$

dijet

summary

$$\sigma = \frac{N_{\text{candidates}} - N_{\text{background}}}{\epsilon_{\text{cosmic}} \epsilon_{\text{fsr}} \epsilon_{\text{ee}} \mathcal{L}_{\text{eff}}}$$

$$N_{\text{candidates}} = 16^{+5.1}_{-3.2} \text{ (stat)} \quad \epsilon_{\text{fsr}} = 0.79 \pm 0.05 \text{ (sys)}$$

$$N_{\text{background}} = 2.1^{+0.7}_{-0.3} \text{ (sys)} \quad \epsilon_{\text{cosmic}} = 0.93 \pm 0.03 \text{ (sys)}$$

$$\mathcal{L}_{\text{eff}} = 46 \pm 3 \text{ (sys) pb}^{-1} \quad \epsilon_{\text{ee}} = 0.26 \pm 0.03 \text{ (sys)}$$

$$\sigma_{\text{MEASURED}} = 1.6^{+0.5}_{-0.3} \text{ (stat)} \pm 0.3 \text{ (sys) pb}$$

**Poisson probability of $2.8 \rightarrow 16 = 5.0 \times 10^{-8}$
Corresponds to 5.4σ “observation”**

Agrees with LPAIR theory: $\sigma_{\text{LPAIR}} = 1.711 \pm 0.008 \text{ pb}$



ee Numbers



introduction

outline

DØ plans

CDF inclusive

intro

results

CDF exclusive

intro

e^+e^-

$\gamma\gamma$

dijet

summary

Fakes: $0.0^{+0.1}_{-0.0}$ events

Cosmic: negligible

Exclusive: $0.0^{+0.3}_{-0.0}$ events

Dissociation: 2.1 ± 0.3 events





Total: $2.1^{+0.7}_{-0.3}$ events



ee Plots



introduction
outline
DØ plans
CDF inclusive
 intro
 results
CDF exclusive
 intro
 e^+e^-
 $\gamma\gamma$
 dijet
summary





Dijet Plots



introduction

outline

DØ plans

CDF inclusive

intro

results

CDF exclusive

intro

e^+e^-

$\gamma\gamma$

dijet

summary


DATASETS:

Exclusive / Inclusive fraction: 310 pb^{-1}

Heavy Flavor fraction: 200 pb^{-1}


ExHuME

CDF Run II Preliminary



DPEMC

CDF Run II Preliminary










Dijet Plots




- introduction
- outline
- DØ plans
- CDF inclusive
 - intro
 - results
- CDF exclusive
 - intro
 - e^+e^-
 - $\gamma\gamma$
 - dijet
- summary




CDF Run II Preliminary




CDF Run II Preliminary



CDF Run II Preliminary



CDF Run II Preliminary





Dijet Cross Section



introduction

outline

DØ plans

CDF inclusive

intro

results

CDF exclusive

intro

e^+e^-

$\gamma\gamma$

dijet

summary

CDF Run II Preliminary

$E_T^{\text{jet1,2}} > E_T^{\text{min}}$ GeV, $|\eta^{\text{jet1,2}}| < 2.5$, $3.6 < \eta_{\text{gap}} < 5.9$, $0.03 < \xi_{\bar{p}} < 0.08$, all $t_{\bar{p}}$

E_T^{min}	σ_{incl}	σ_{excl}	$\sigma_{\text{excl}}/\sigma_{\text{incl}}$
10	$14.5 \pm 0.1^{+9.8}_{-6.9}$ nb	$1.10 \pm 0.04^{+1.29}_{-0.54}$ nb	$7.6 \pm 0.3^{+2.9}_{-1.2}$ %
15	$1.43 \pm 0.02^{+0.89}_{-0.62}$ nb	$112 \pm 7^{+84}_{-49}$ pb	$7.8 \pm 0.5^{+3.2}_{-1.2}$ %
20	$267 \pm 6^{+166}_{-110}$ pb	$15.7 \pm 2.0^{+15.5}_{-9.6}$ pb	$5.9 \pm 0.8^{+3.0}_{-2.1}$ %
25	$76.0 \pm 2.7^{+37.0}_{-28.6}$ pb	$4.84 \pm 0.96^{+4.11}_{-3.28}$ pb	$6.4 \pm 1.3^{+4.6}_{-3.9}$ %
35	$14.6 \pm 1.2^{+5.3}_{-5.2}$ pb	$1.37 \pm 0.49^{+1.08}_{-1.01}$ pb	$9.3 \pm 3.4^{+6.9}_{-6.6}$ %

(stat. + syst. errors)